

UDK 512.815.1+512.815.6+512.816.1+512.816.2

MSC 14L24+22C05

O. G. Styrť

The existence of a semialgebraic continuous factorization map for some compact linear groups

It is proved that each of compact linear groups of one special type admits a semialgebraic continuous factorization map onto a real vector space.

Key words: Lie group, factorization map of an action, semialgebraic map.

§ 1. Introduction

In this paper, we prove that each of compact linear groups of one certain type admits a semialgebraic continuous factorization map onto a real vector space. This problem arose from the question when the topological quotient of a compact linear group is homeomorphic to a vector space that was researched in [1, 2, 3, 4, 5].

For convenience, denote by $\mathcal{P}_{\mathbb{R}}$ (resp. by $\mathcal{P}_{\mathbb{C}}$) the class of all finite-dimensional Euclidean (resp. Hermitian) spaces. Let $\mathbb{F}$ be one of the fields $\mathbb{R}$ and $\mathbb{C}$. We will write $\mathbb{F}^k$ ($k \geq 0$) for the space $\mathbb{F}^k \in \mathcal{P}_{\mathbb{F}}$ whose standard basis is orthonormal. For any spaces $W_1, W_2 \in \mathcal{P}_{\mathbb{F}}$, we will write $W_1 \oplus W_2$ for their *orthogonal* direct sum $W_1 \oplus W_2 \in \mathcal{P}_{\mathbb{F}}$ (unless they are linearly independent subspaces of the same space). For any space $W \in \mathcal{P}_{\mathbb{F}}$, denote by $S(W)$ the real subspace $\{A \in \text{End}_{\mathbb{F}}(W) : A = A^*\} \subset \text{End}_{\mathbb{F}}(W)$, by $\mathbf{O}(W)$ the compact Lie group $\{C \in \mathbf{GL}_{\mathbb{F}}(W) : CC^* = E\} \subset \mathbf{GL}_{\mathbb{F}}(W)$, and by $\mathbf{SO}(W)$ its compact subgroup $\mathbf{O}(W) \cap \mathbf{SL}_{\mathbb{F}}(W)$.

The following theorem is the main result of the paper and will be proved in § 3.

Theorem 1.1. *Consider a number $k \in \{1, 2\}$, a space $W \in \mathcal{P}_{\mathbb{F}}$ such that $\dim_{\mathbb{F}} W \geq 2-k$, and the representation*

$$\mathbf{O}(W) : (S(W)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W), \quad C : (A + \mathbb{R}E, B) \rightarrow (CAC^{-1} + \mathbb{R}E, CB). \quad (1.1)$$

- 1) If $k = 1$, then there exist a real vector space V and, in terms of the representation $\mathbf{O}(W): V \oplus \mathbb{F}$, $C: (v, \lambda) \rightarrow (v, \det_{\mathbb{F}} C \cdot \lambda)$, a semialgebraic continuous $(\mathbf{O}(W))$ -equivariant surjective map $(S(W)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W) \rightarrow V \oplus \mathbb{F}$ whose fibres coincide with the $(\mathbf{SO}(W))$ -orbits.
- 2) If $k = 2$, then there exist a real vector space V and a semialgebraic continuous surjective map $(S(W)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W) \rightarrow V$ whose fibres coincide with the $(\mathbf{O}(W))$ -orbits.

§ 2. Auxiliary facts

Consider a space $W \in \mathcal{P}_{\mathbb{F}}$ and the subsets $S_+(W) := \{A \in S(W) : A \geq 0\} \subset S(W)$ and $S_0(W) := (S_+(W)) \setminus (\mathbf{GL}_{\mathbb{F}}(W)) \subset S_+(W) \subset S(W)$.

Set $M(W) := \{(A, \lambda) \in S_+(W) \times \mathbb{F} : \det_{\mathbb{F}} A = |\lambda|^2\} \subset S(W) \oplus \mathbb{F}$.

The following statement is well-known.

Statement 2.1. *The fibres of the map $\pi: \text{End}_{\mathbb{F}}(W) \rightarrow S(W) \oplus \mathbb{F}$, $X \rightarrow (XX^*, \det_{\mathbb{F}} X)$ coincide with the orbits of the action*

$$\mathbf{SO}(W): \text{End}_{\mathbb{F}}(W), C: X \rightarrow XC^{-1}, \quad (2.1)$$

and $\pi(\text{End}_{\mathbb{F}}(W)) = M(W) \subset S(W) \oplus \mathbb{F}$.

Lemma 2.1. *The map $M(W) \rightarrow (S(W)/(\mathbb{R}E)) \oplus \mathbb{F}$, $(A, \lambda) \rightarrow (A + \mathbb{R}E, \lambda)$ is a bijection.*

□ See Lemma 5.1 in [3]. ■

Corollary 2.1. *The map $S_0(W) \rightarrow S(W)/(\mathbb{R}E)$, $A \rightarrow A + \mathbb{R}E$ is a bijection.*

Corollary 2.2. *The map $\text{End}_{\mathbb{F}}(W) \rightarrow (S(W)/(\mathbb{R}E)) \oplus \mathbb{F}$, $X \rightarrow (XX^* + \mathbb{R}E, \det_{\mathbb{F}} X)$ is surjective, and its fibres coincide with the orbits of the action (2.1).*

□ Follows from Statement 2.1 and Lemma 2.1. ■

Now consider arbitrary spaces $W, W_1 \in \mathcal{P}_{\mathbb{F}}$ over the field $\mathbb{F}$.

Statement 2.2. *The fibres of the map $\pi: \text{Hom}_{\mathbb{F}}(W_1, W) \rightarrow S(W)$, $X \rightarrow XX^*$ coincide with the orbits of the action*

$$\mathbf{O}(W_1): \text{Hom}_{\mathbb{F}}(W_1, W), C_1: X \rightarrow XC_1^{-1}, \quad (2.2)$$

and $\pi(\text{Hom}_{\mathbb{F}}(W_1, W)) = \{A \in S_+(W) : \text{rk}_{\mathbb{F}} A \leq \dim_{\mathbb{F}} W_1\} \subset S(W)$. In particular, in the case $\dim_{\mathbb{F}} W_1 = \dim_{\mathbb{F}} W - 1$, we have $\pi(\text{Hom}_{\mathbb{F}}(W_1, W)) = S_0(W) \subset S(W)$.

We omit the proof since it is clear.

Corollary 2.3. *Suppose that $\dim_{\mathbb{F}} W_1 = \dim_{\mathbb{F}} W - 1$. Then the map $\text{Hom}_{\mathbb{F}}(W_1, W) \rightarrow S(W)/(\mathbb{R}E)$, $X \rightarrow XX^* + \mathbb{R}E$ is surjective, and its fibres coincide with the orbits of the action (2.2).*

□ Follows from Statement 2.2 and Corollary 2.1. ■

§ 3. Proof of the main result

In this section, we will prove Theorem 1.1.

First of all, let us prove the claim for a pair (k, W) ($k \in \{1, 2\}$, $W \in \mathcal{P}_{\mathbb{F}}$, $\dim_{\mathbb{F}} W = 2 - k$).

If $k = 2$ and $W = 0$, then the space of the representation (1.1) is trivial. If $k = 1$ and $\dim_{\mathbb{F}} W = 1$, then the representation (1.1) is isomorphic to the tautological representation of the group $\mathbf{O}(W)$ and, consequently, to the representation $\mathbf{O}(W): \mathbb{F}, C: \lambda \rightarrow \det_{\mathbb{F}} C \cdot \lambda$, and the subgroup $\mathbf{SO}(W) \subset \mathbf{O}(W)$ is trivial.

This completely proves the claim for a pair (k, W) ($k \in \{1, 2\}$, $W \in \mathcal{P}_{\mathbb{F}}$, $\dim_{\mathbb{F}} W = 2 - k$).

Take an arbitrary pair (k, W) ($k \in \{1, 2\}$, $W \in \mathcal{P}_{\mathbb{F}}$, $n := \dim_{\mathbb{F}} W > 2 - k$) and assume that the claim of Theorem 1.1 holds for any pair (k, W_1) ($W_1 \in \mathcal{P}_{\mathbb{F}}$, $\dim_{\mathbb{F}} W_1 = n - 1$). We will now prove it for the pair (k, W) .

Since $n > 2 - k \geq 0$, there exists a space $W_1 \in \mathcal{P}_{\mathbb{F}}$ such that $\dim_{\mathbb{F}} W_1 = n - 1$. Consider the embedding $R: \mathbf{O}(W_1) \hookrightarrow \mathbf{O}(W_1 \oplus \mathbb{F}^k)$, $R(C_1): (x_1, x_2) \rightarrow (C_1 x_1, x_2)$. The representations

$$\mathbf{O}(W_1): S(W_1 \oplus \mathbb{F}^k), C_1: A \rightarrow R(C_1)AR(C_1^{-1}) \quad (3.1)$$

and $\mathbf{O}(W_1): S(W_1) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1) \oplus S(\mathbb{F}^k)$, $C_1: (A_1, B, A_2) \rightarrow (C_1 A_1 C_1^{-1}, C_1 B, A_2)$ are isomorphic via the $(\mathbf{O}(W_1))$ -equivariant $\mathbb{R}$ -linear isomorphism

$$\begin{aligned} \theta: S(W_1) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1) \oplus S(\mathbb{F}^k) &\rightarrow S(W_1 \oplus \mathbb{F}^k), \\ \theta(A_1, B, A_2): (x_1, x_2) &\rightarrow (A_1 x_1 + B x_2, B^* x_1 + A_2 x_2). \end{aligned}$$

Therefore, the representation (3.1) is isomorphic to the direct sum of some trivial real representation of the group $\mathbf{O}(W_1)$ and the representation

$$\mathbf{O}(W_1): (S(W_1)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1), C_1: (A_1 + \mathbb{R}E, B) \rightarrow (C_1 A_1 C_1^{-1} + \mathbb{R}E, C_1 B)$$

with trivial fixed point subspace. Thus, there exist a real vector space V' and an isomorphism $\varphi: S(W_1 \oplus \mathbb{F}^k)/(\mathbb{R}E) \rightarrow V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1)$ of the representations

$$\begin{aligned} \mathbf{O}(W_1): S(W_1 \oplus \mathbb{F}^k)/(\mathbb{R}E), C_1: A + \mathbb{R}E &\rightarrow R(C_1)AR(C_1^{-1}) + \mathbb{R}E; \\ \mathbf{O}(W_1): V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1), \\ C_1: (v', A_1 + \mathbb{R}E, B) &\rightarrow (v', C_1 A_1 C_1^{-1} + \mathbb{R}E, C_1 B). \end{aligned}$$

In terms of the representations

$$\begin{aligned} \mathbf{O}(W) \times \mathbf{O}(W_1): \text{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W), (C, C_1): Z &\rightarrow CZR(C_1^{-1}); \\ \mathbf{O}(W) \times \mathbf{O}(W_1): (S(W)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W), \\ (C, C_1): (A + \mathbb{R}E, B) &\rightarrow (CAC^{-1} + \mathbb{R}E, CB); \\ \mathbf{O}(W) \times \mathbf{O}(W_1): V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1), \\ (C, C_1): (v', A_1 + \mathbb{R}E, B) &\rightarrow (v', C_1 A_1 C_1^{-1} + \mathbb{R}E, C_1 B), \end{aligned}$$

the maps

$$\begin{aligned}\pi_0: \operatorname{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W) &\rightarrow (S(W)/(\mathbb{R}E)) \oplus \operatorname{Hom}_{\mathbb{F}}(\mathbb{F}^k, W), \\ Z &\rightarrow ((Z|_{W_1})(Z|_{W_1})^* + \mathbb{R}E, Z|_{\mathbb{F}^k}); \\ \psi: \operatorname{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W) &\rightarrow V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \operatorname{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1), \quad Z \rightarrow \varphi(Z^*Z + \mathbb{R}E)\end{aligned}$$

are $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant. By Corollary 2.3, the map π_0 is surjective, and its fibres coincide with the $(\mathbf{O}(W_1))$ -orbits.

Case 1). $k = 1$.

Consider the representation

$$\begin{aligned}\mathbf{O}(W) \times \mathbf{O}(W_1): V' \oplus (S(W_1)/(\mathbb{R}E)) &\oplus \operatorname{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1) \oplus \mathbb{F}, \\ (C, C_1): (v', A_1 + \mathbb{R}E, B, \mu) &\rightarrow (v', C_1 A_1 C_1^{-1} + \mathbb{R}E, C_1 B, \det_{\mathbb{F}} C \cdot \det_{\mathbb{F}} C_1^{-1} \cdot \mu).\end{aligned}$$

Since $\dim_{\mathbb{F}} W_1 = n - 1$, there exist a real vector space V_1 and, in terms of the representation $\mathbf{O}(W) \times \mathbf{O}(W_1): V' \oplus V_1 \oplus \mathbb{F}^2$, $(C, C_1): (v', v_1, \lambda, \mu) \rightarrow (v', v_1, \det_{\mathbb{F}} C_1 \cdot \lambda, \det_{\mathbb{F}} C \cdot \det_{\mathbb{F}} C_1^{-1} \cdot \mu)$, a semialgebraic continuous $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant surjective map

$$\pi_1: V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \operatorname{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1) \oplus \mathbb{F} \rightarrow V' \oplus V_1 \oplus \mathbb{F}^2$$

whose fibres coincide with the $(\mathbf{SO}(W_1))$ -orbits. Since $\dim_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k) = (n - 1) + k = n = \dim_{\mathbb{F}} W$, we can identify W and $W_1 \oplus \mathbb{F}^k$ as spaces of the class $\mathcal{P}_{\mathbb{F}}$. The map

$$\begin{aligned}\tilde{\psi}: \operatorname{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W) &\rightarrow V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \operatorname{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1) \oplus \mathbb{F}, \\ Z &\rightarrow (\psi(Z), \det_{\mathbb{F}} Z) = (\varphi(Z^*Z + \mathbb{R}E), \det_{\mathbb{F}} Z)\end{aligned}$$

is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant. Also, by Corollary 2.2, the map $\tilde{\psi}$ is surjective, and its fibres coincide with the $(\mathbf{SO}(W))$ -orbits. Hence, the map $\pi_1 \circ \tilde{\psi}: \operatorname{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W) \rightarrow V' \oplus V_1 \oplus \mathbb{F}^2$ is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant and surjective, and its fibres coincide with the $(\mathbf{SO}(W) \times \mathbf{SO}(W_1))$ -orbits. If we consider the representation

$$\mathbf{O}(W) \times \mathbf{O}(W_1): V' \oplus V_1 \oplus \mathbb{R} \oplus \mathbb{F}, \quad (C, C_1): (v', v_1, t, \lambda) \rightarrow (v', v_1, t, \det_{\mathbb{F}} C \cdot \lambda),$$

then the map $\gamma: V' \oplus V_1 \oplus \mathbb{F}^2 \rightarrow V' \oplus V_1 \oplus \mathbb{R} \oplus \mathbb{F}$, $(v', v_1, \lambda, \mu) \rightarrow (v', v_1, |\lambda|^2 - |\mu|^2, \lambda\mu)$ is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant. Denote by $\mathbb{T}$ the multiplicative group $\{c \in \mathbb{F}: |c| = 1\}$. We have $n > 2 - k = 1$, $\dim_{\mathbb{F}} W_1 = n - 1 > 0$. Consequently,

$$\{\det_{\mathbb{F}} C_1: C_1 \in \mathbf{O}(W_1)\} = \mathbb{T}. \quad (3.2)$$

One can easily see that the map $\mathbb{F}^2 \rightarrow \mathbb{R} \oplus \mathbb{F}$, $(\lambda, \mu) \rightarrow (|\lambda|^2 - |\mu|^2, \lambda\mu)$ is surjective, and its fibres coincide with the orbits of the representation $\mathbb{T}: \mathbb{F}^2, c: (\lambda, \mu) \rightarrow (c\lambda, c^{-1}\mu)$. By (3.2), the map γ is surjective, and its fibres coincide with the $(\mathbf{O}(W_1))$ -orbits. It follows from above

that the map $\gamma \circ \pi_1 \circ \tilde{\psi}: \text{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W) \rightarrow V' \oplus V_1 \oplus \mathbb{R} \oplus \mathbb{F}$ is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant and surjective, and its fibres coincide with the $(\mathbf{SO}(W) \times \mathbf{O}(W_1))$ -orbits.

Recall that the map π_0 is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant and surjective, and its fibres coincide with the $(\mathbf{O}(W_1))$ -orbits. Therefore, there exists an $(\mathbf{O}(W))$ -equivariant surjective map $\pi: (S(W)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W) \rightarrow V' \oplus V_1 \oplus \mathbb{R} \oplus \mathbb{F}$ satisfying $\pi \circ \pi_0 \equiv \gamma \circ \pi_1 \circ \tilde{\psi}$ whose fibres coincide with the $(\mathbf{SO}(W))$ -orbits. Since the maps $\gamma, \pi_1, \tilde{\psi}, \pi_0$ are semialgebraic, continuous, and surjective, so is the map π .

This completely proves the claim in the case $k = 1$.

Case 2). $k = 2$.

Since $\dim_{\mathbb{F}} W_1 = n - 1$, there exist a real vector space V_1 and a semialgebraic continuous surjective map $\pi_1: V' \oplus (S(W_1)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W_1) \rightarrow V' \oplus V_1$ whose fibres coincide with the $(\mathbf{O}(W_1))$ -orbits. Further, $\dim_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k) = (n - 1) + k = n + 1 = \dim_{\mathbb{F}} W + 1$, and, by Corollary 2.3, the map ψ is surjective, and its fibres coincide with the $(\mathbf{O}(W))$ -orbits. Since the map ψ is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant, the map $\pi_1 \circ \psi: \text{Hom}_{\mathbb{F}}(W_1 \oplus \mathbb{F}^k, W) \rightarrow V' \oplus V_1$ is surjective, and its fibres coincide with the $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -orbits. Recall that the map π_0 is $(\mathbf{O}(W) \times \mathbf{O}(W_1))$ -equivariant and surjective, and its fibres coincide with the $(\mathbf{O}(W_1))$ -orbits. Hence, there exists a surjective map $\pi: (S(W)/(\mathbb{R}E)) \oplus \text{Hom}_{\mathbb{F}}(\mathbb{F}^k, W) \rightarrow V' \oplus V_1$ satisfying $\pi \circ \pi_0 \equiv \pi_1 \circ \psi$ whose fibres coincide with the $(\mathbf{O}(W))$ -orbits. Since the maps π_1, ψ, π_0 are semialgebraic, continuous, and surjective, so is the map π .

This completely proves the claim in the case $k = 2$.

Thus, we have completely proved Theorem 1.1 (using mathematical induction on $\dim_{\mathbb{F}} W$ separately for each number $k \in \{1, 2\}$).

References

- [1] M. A. Mikhailova, *On the quotient space modulo the action of a finite group generated by pseudoreflections*, Mathematics of the USSR-Izvestiya, 1985, vol. 24, 1, 99—119.
- [2] O. G. Styr, *On the orbit space of a compact linear Lie group with commutative connected component*, Tr. Mosk. Mat. O-va, 2009, vol. 70, 235—287 (Russian).
- [3] O. G. Styr, *On the orbit space of a three-dimensional compact linear Lie group*, Izv. RAN, Ser. math., 2011, vol. 75, 4, 165—188 (Russian).
- [4] O. G. Styr, *On the orbit space of an irreducible representation of a special unitary group*, Tr. Mosk. Mat. O-va, 2013, vol. 74, 1, 175—199 (Russian).
- [5] O. G. Styr, *On the orbit spaces of irreducible representations of simple compact Lie groups of types B, C, and D*, J. Algebra, 2014, vol. 415, 137—161.